

GLP-1 Receptor Agonist

Medications used in the treatment of Type 2 Diabetes and Weight Loss

Hold for day of procedure:

Byetta® (exenatide)
Saxenda® (liraglutide)
Victoza® (liraglutide)
Adlyxin® (lixisenatide)
Rybelsus® (semaglutide)

Xultophy®
(liraglutide/insulin
degludec)

Soliqua®
(lixisenatide/insulin)

Inform patients that if they hold their medicine as directed and they have GI symptoms the day of procedure (nausea, bloating, abdominal pain, etc.) the anesthesiologist may elect to cancel the procedure due to risk of aspiration.


Hold for 7 days:

Trulicity® (dulaglutide)
Bydureon® (exenatide)
Ozempic® (semaglutide)
Wegovy® (semaglutide)
Mounjaro® (tirzepatide)
Zepbound™ (tirzepatide)

**Rule of thumb: If the GLP-1 is prescribed for daily use, hold day of procedure.
If the GLP-1 is prescribed weekly, hold for 7 days.**

SGLT-2 Inhibitors

Medications used in the treatment of Diabetes Mellitus Type 2, Heart Failure and Chronic Kidney Disease



Examples of SGLT-2 Inhibitors: **Brenzavvy**® (bexagliflozin), **Invokana**® (canagliflozin), **Invokamet**® (canagliflozin/metformin), **Farxiga**® (dapagliflozin), **Xigduo XR**® (dapagliflozin/metformin), **Qtern**® (dapagliflozin/saxagliptin), **Jardiance**® (empagliflozin), **Glyxambi**® (empagliflozin/linagliptin), **Synjardy**®/**Synjardy XR**® (empagliflozin/metformin), **Trijardy**®/**Trijardy XR**® (empagliflozin/metformin/linagliptin), **Steglatro**® (ertugliflozin), **Inpefa**® (Sotagliflozin), **Segluromet**® (ertugliflozin/metformin), **Steglujan**® (ertugliflozin/sitagliptin)