

Weight Loss/Diabetes Medication and Procedures

How long do I hold my medicine before my procedure:

	Minor surgeries/procedures	Intermediate/high risk surgeries
bexagliflozin (Brenzavvy®)	Day of procedure	3 days prior
Canagliflozin (Invokana®, Invokamet®)	Day of procedure	3 days prior
dapagliflozin (Farxiga®, Xigduo XR®, Qtern®)	Day of procedure	3 days prior
empagliflozin (Jardiance®, Glyxambi®, Synjardy®, Trijardy®)	Day of procedure	3 days prior
ertugliflozin (Steglatro®, Segluromet®, Steglujan®)	Day of procedure	4 days prior
Sotagliflozin (Inpefa®)	Day of procedure	3 days prior
Dulaglutide (Trulicity®)	7 days prior (skip a dose)	7 days prior (skip a dose)
Exenatide ER (Bydureon bcise®)	7 days prior (skip a dose)	7 days prior (skip a dose)
Semaglutide injection (Ozempic®, Wegovy®)	7 days prior (skip a dose)	7 days prior (skip a dose)
Tirzepatide (Mounjaro®, Zebound®)	7 days prior (skip a dose)	7 days prior (skip a dose)
Exenatide IR (Byetta®)	Day of procedure	Day of procedure
Liraglutide (Victoza®, Saxenda®, Xultophy®)	Day of procedure	Day of procedure
Lixisenatide (Adlyxin®, Soliqua®)	Day of procedure	Day of procedure
Semaglutide Oral (Rybelsus®)	Day of procedure	Day of procedure

Minor surgeries/procedures: TEE, cardioversions, Pacemakers, Defibrillators, Loop Recorders, Heart caths, SVT ablations, EGDs, colonoscopies, etc). Schedule patients earlier in the day (AM preferred) to avoid unnecessary NPO time

Intermediate/High Risk surgeries/procedures: Afib/flutter ablations, VT ablations, Mitral Clips, TAVRs, etc.